

Problem Set 3

March 3, 2019.

1. Let X be a k -dimensional analytic subset of a manifold M , and let, for every $d \in \mathbb{N}$,

$$X^{(d)} = \{x \in X : \dim_x X = d\}.$$

Prove the following:

- (a) $X^{(k)}$ is an analytic subset of X (in particular, closed!).
 (b) For every $d \in \mathbb{N}$, $\overline{X^{(d)}}$ is analytic of pure dimension d . Moreover,

$$X = X^{(k)} \cup X^{(k-1)} \cup \dots \cup X^{(0)} = X^{(k)} \cup \overline{X^{(k-1)}} \cup \dots \cup \overline{X^{(0)}}.$$

- (c) For every $d = 1, \dots, k$, $\bigcup_{i < d} X^{(i)}$ is open in X .
 (d) For every $d = 1, \dots, k$, $X \setminus \bigcup_{i < d} X^{(i)}$ is analytic in X .

2. Let X and $X^{(d)}$ be as above. Give, as explicit as possible, a characterization of $X^{(d)}$ ($d = 0, \dots, k$) in terms of components of $\text{Reg}(X)$.

3. Prove Proposition 8.8 and Remark 8.13(1) from Lecture Notes.

4. (a) [Hartshorne, Ex.II.1.2(c)] Show that a sequence of sheaves on a topological space X is exact iff, for every $\xi \in X$, the corresponding sequence of stalks at ξ is exact.
 (b) Use part (a) to conclude that a morphism of sheaves $\alpha : \mathcal{F} \rightarrow \mathcal{G}$ is injective (resp. surjective) iff α_ξ is so for every $\xi \in X$.

5. [Hartshorne, Ex.II.1.3]

- (a) Let $\alpha : \mathcal{F} \rightarrow \mathcal{G}$ be a morphism of sheaves on X . Show that α is surjective iff the following condition holds: for every open $U \subset X$, and for every $t \in \mathcal{G}(U)$, there is an open covering $\{U_i\}_i$ of U , and sections $s_i \in \mathcal{F}(U_i)$, such that $\alpha(U_i)(s_i) = t|_{U_i}$ for all i .
 (b) Give an example of a surjective morphism of sheaves $\alpha : \mathcal{F} \rightarrow \mathcal{G}$, and an open set U , such that $\alpha(U) : \mathcal{F}(U) \rightarrow \mathcal{G}(U)$ is not surjective.

6. [Hartshorne, Ex.II.1.19] (*Extending a sheaf by zero*) Let X be a topological space, let Z be a closed subset, let $U = X \setminus Z$, and let $i : Z \rightarrow X$ and $j : U \rightarrow X$ be the inclusions.

- (a) Let \mathcal{F} be a sheaf on Z . Show that the stalk $(i_*\mathcal{F})_x$ of the direct image sheaf on X is \mathcal{F}_x if $x \in Z$, and 0 if $x \notin Z$.
 (b) Now let \mathcal{F} be a sheaf on U . Let $j_!(\mathcal{F})$ be the sheaf on X obtained as a sheafification of the presheaf $\{V \mapsto \mathcal{F}(V) \text{ if } V \subset U, V \mapsto 0 \text{ otherwise}\}$. Show that the stalk $(j_!(\mathcal{F}))_x$ is equal to \mathcal{F}_x if $x \in U$, and 0 if $x \notin U$, and show that $j_!(\mathcal{F})$ is the only sheaf on X which has this property, and whose restriction to U is \mathcal{F} .

7. Prove Lemma 8.20 and Corollary 8.28 from Lecture Notes.